

Understanding Pica: An Exploration of an Uncommon Eating Disorder

Ms. Lina Sarkar

Assistant Professor, Department of Psychology,
Swami Vivekananda University

Introduction

Pica is an eating disorder characterized by the compulsive consumption of non-food items that lack nutritional value. While it is common for young children to explore their environment by putting various objects in their mouths, pica extends beyond this natural curiosity. In some cases, it can lead to significant health issues, particularly when hazardous substances are ingested.

Pica is an eating disorder where individuals crave or consume non-foods items which can be natural, biological, or man-made. The name comes from the Latin word for magpie—a bird known for eating a wide variety of things. According to the DSM-5 to be diagnosed with pica, the behavior must last for more than a month and be considered inappropriate for the person's developmental stage. It should not be part of a cultural practice and be serious enough to require clinical attention. Pica poisoning can occur, especially in children, affecting their physical and mental development. It can also cause serious health issues like intestinal blockages, nutritional deficiencies and infections from parasites.



Classification of Pica

Different types of Pica include-

- Geophagia: Including dirt, soil, clay or sand eating.
- Pagophagia: Including eating ice, frosty, or uncontrollably drinking cooled beverage –frequently observed in individuals with anemia.
- Amylophagia: Including uncooked rice or pasta, laundry starch etc.

Who is Affected by Pica?

Pica can affect individuals of any age, but it is most prevalent in three groups:


- Young Children: Particularly those under the age of six, who may engage in pica behaviors as part of normal development.
- Pregnant Individuals: Hormonal changes can increase cravings for non-food items.
- Individuals with Mental Health Conditions: Those diagnosed with autism spectrum disorder, intellectual disabilities, or schizophrenia are at a higher risk.

Understanding Pica in Young Children: Causes, Risks, and Management Strategies-

Pica is characterized by the persistent consumption of non-eatable substances, last at least one month, which is not suitable for developmental stages. It is conventional among young children and individuals who have developmental disabilities.

Causes of Pica in Young Children

- Young children understand their environment using their senses, including taste, which perceives non-food items.
- Sometimes children consume nonfood items to compensate nutrients deficiencies such as iron or zinc.
- Stress, anxiety, or mental health conditions may cause children to eat non-food items as a coping mechanism.
- In some cultures, eating certain non-food items is considered normal or traditional.



Causes of Pica in Young Children

Risks and Management Strategies:

Addressing underlying causes, provide appropriate supervision, and providing appropriate medical and psychological interventions from professionals involves management and treatment for pica behavior. Nutritional deficiencies should be corrected, and behavioral therapy can be used to discourage pica behavior. Educating parents and creating a safe environment can help prevent eating non-food. If pica is suspected, it is crucial to seek professional medical advice for proper diagnosis and treatment.

Understanding Pica in Pregnant Individuals: Causes, Risks, and Management Strategies

Pica is a condition characterized by the persistent consumption of non-dietary substances for at least 1 month, which is not suitable to the developmental level of the individual. This condition isn't only prevalent among children but also among pregnant individuals.

Causes of Pica in Pregnant Individuals

Insufficiencies in nutrients can cause pica behavior, as can the body, because the body may try to obtain these nutrients through non-eatable items. Due to certain belief system and cultural practices may normalize the intake of non-eatable items during pregnancy. Stress, anxiety, or underlying mental health conditions can contribute to pica as a coping mechanism. Fluctuations in pregnancy can cause unusual cravings, including for non-eatable items. Some pregnant woman may develop a liking for the smell, texture or taste of non-eatable items.

CAUSE	% (PERCENTAGE)
Hormonal Changes	20%
Sensory Appeal	20%
Nutritional Deficiencies	20%
Cultural and Societal Factors	15%
Psychological Factors	25%

Percentages of each cause of pica in pregnant individual

Risks Associated with Pica in Pregnancy

Consuming non-eatable items can interfere with the exploitation of essential nutrients, leading to deficiencies. To eat materials such as soil, clay or chalk can expose a person to harmful toxins and pathogens. Dental problem occurs due to chewing on hard non-food items, including tooth breakage. Eating indigestible items can cause barrier in the digestive system.

Management Strategies

- Education is required for pregnant women about the risks related to pica and to motivate them for adopt healthy eating habits.
- For monitoring and addressing any health-related problems arising from pica, regular check-ups are required.
- Identify nutritional deficit.
- To reduce stress, anxiety and other psychological factors provide them counselling.

Causes and Management Strategies- Understanding pica in individuals with mental health conditions:

Understanding pica in individuals with mental health conditions includes various factors that contribute to this behavior, as well as strategies for management. A brief discussion is provided below-

Causes of Pica in Individuals with Mental Health Conditions

Deficiencies in nutrients including iron, zinc, or calcium may cause people with mental health problems to crave non-food items. Obsessive-compulsive disorder (OCD), autism spectrum disorder (ASD), schizophrenia, & intellectual disability can contribute to pica. It can serve as a coping mechanism for stress, anxiety, or emotional distress. Some people with mental health conditions may have heightened sensory experiences, leading them to seek out specific textures or tastes that non-food items provide. Staying in an environment where non-food items are accessible or normalized, or social factors that strengthen pica behavior,

may affect individuals with mental health conditions.

Causes of Pica in Individuals with Mental Health Conditions	Percentage(%)
Nutritional Deficiencies	25%
Psychological Factors	35%
Sensory Sensitivity	15%
Environmental and Social Factors	25%

Risks Associated with Pica

Intake non-food items can lead to bowel obstruction, dental problems, poisoning and other serious physical health complications. Pica can deepen existing nutritional deficiencies or lead to imbalances, further impacting overall health. Consuming toxic or sharp substances can result in physical harm, such as cuts, wounds or poisoning.

Management Strategies

- Identifying underlying medical conditions, nutritional insufficiencies, or psychiatric disorders that lead to pica requires comprehensive assessment by healthcare professionals.
- To address the underlying causes of pica and reduce the urge to eat non-food items Cognitive- behavioral therapy (CBT) and behavior modification techniques can be effective.
- Provide education to caregivers and providing support to families to understand and manage pica behaviors effectively.
- Identifying nutritional deficiencies through dietary adjustments or supplementation under medical supervision.
- Limiting access to non-eatable items and creating a safe living environment to reduce opportunities for pica.

Prevalence and Detection

The specific prevalence of pica is difficult to define due to inconsistent definitions in research and the tendency of healthcare providers to overlook it. Symptoms often go

undisclosed, especially in young children, leading to under diagnosis. Moreover, behaviors in children may not be classified as pica, as they typically resolve on their own.

Health Implications of Pica

Pica can lead to various health complications depending on the types of non-food items intake. While some behaviors, such as eating ice, may be benign, others can result in serious issues:

- Chewing on hard objects can harm teeth (Dental Damage)
- Ingesting hair or large non-food items can cause intestinal obstruction.
- Consuming dirt or feces can lead to infections.

Psychological Impact

Due to feelings of shame or embarrassment, individuals with pica may be reluctant to seek help. This psychological burden complicates the condition, making it essential for healthcare providers to approach the issue sensitively.

Symptoms of Pica

The primary symptom of pica is the compulsive consumption of non-food items. Individuals may develop preferences for specific substances, ranging from chalk and dirt to more unusual items like paint chips and laundry starch.

Factors behind Pica

While the specific cause of pica remains unclear, several risk factors have been identified:

- Cultural Practices: Some cultures accept certain types of pica, such as eating clay or soil for nutritional deficiencies.
- Nutritional Deficiencies: Deficiencies in iron, calcium, or zinc may lead to pica behaviors.
- Psychological Factor: Stress, anxiety, or childhood trauma may contribute to the development of pica as a coping mechanism.

Diagnosis and Assessment

Diagnosing pica requires meeting specific criteria:

- **Duration:** The behavior must persist for at least one month.
- **Mental Development:** The individual should possess the cognitive ability to distinguish food from non-food.
- **Exclusion of Cultural Norms:** The behavior should not be explained by social or cultural practices.
- **Missing of Other Disorders:** Pica should not be primarily attributed to other medical or psychological conditions.

Diagnostic Tests.

To identify health complications related to pica, such as infections or electrolyte imbalances, internal damage etc. diagnosis is required, which may involve lab test.

Treatment and Management

Treatment for pica varies depending on individual situation. In children and pregnant individuals, pica often resolves on its own. For others, therapy is crucial. Therapeutic options may include:

- **Behavioral Therapy:** Teaching coping strategies to replace pica behaviors.
- **Aversive Conditioning:** Encouraging avoidance of non-food items through mild negative reinforcement.

Living with Pica

For those affected, maintaining open communication with healthcare providers is vital. Discussing pica candidly can facilitate better understanding and treatment. Additionally, removing access to non-food items may help reduce compulsive eating behaviors.

Coping Strategies with Pica

There are many strategies to help people make them empower with pica and try to help them to reduce stress and develop

coping skills when dealing with specific problems. Specific strategies, such as chewing gum, avoiding non-eatable items that you crave can all be helpful coping strategies.

Some examples for parents of autistic children include the following:

- Placing your hand on top of the child's hand so that the child cannot eat the non-eatable items or can blocking their hands. This strategy is helpful to reduce pica for some children.
- Regular time intervals can provide the child with alternatives to eating nonfood items.
- Positive reinforcement, such as toys, can be given when children avoid from pica behaviors. Reward is a proven procedure to help parents change a child's undesirable behaviors.

Conclusion

Pica, though often benign, can lead to serious health consequences if not addressed. Understanding its psychological underpinnings and health implications can empower individuals and healthcare providers to approach the disorder with compassion and effectiveness. As research continues, increased awareness of pica will enhance diagnosis and treatment, supporting individuals in managing this complex condition.

References

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Publishing.
- Bryant-Waugh, R., et al. (2010). Pica in children and adolescents: A review of its clinical presentation, etiology, and treatment. *Journal of Child Psychology and Psychiatry*, 51(5), 561-574.
- Callahan GN. Eating dirt (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3020602/>). *Emerg Infect Dis*.

2003;9(8):1016-1021. Accessed 5/6/2022.

Centers for Disease Control and Prevention. (n.d.). Pica. Retrieved from <https://www.cdc.gov/ncbddd/childdevelopment/nutrition.html>

Leung AKC, Hon KL. Pica: A Common Condition that is Commonly Missed - An Update Review (<https://pubmed.ncbi.nlm.nih.gov/30868957/>). *CurrPediatr Rev.* 2019;15(3):164-169. Accessed 5/6/2022.

Mayo Clinic. (n.d.). Pica. Retrieved from <https://www.mayoclinic.org/diseases-conditions/pica/symptoms-causes/syc-2037627>

Mishori R, McHale C. Pica: an age-old eating disorder that's often missed (<https://pubmed.ncbi.nlm.nih.gov/25198212/>). *J Fam Pract.* 2014;63(7):E1-E4.

National Institutes of Health. (n.d.). Pica. MedlinePlus. Retrieved from [<https://medlineplus.gov/ency/article/001538.htm>] Mayo Clinic. (n.d.). Pica. Retrieved from (<https://www.mayoclinic.org/diseases-conditions/pica/symptoms-causes/syc-2037627>)

Rose, E. A., & Porcerelli, J. H. (2005). Pica: Common but commonly missed. *Journal of the American Academy of Child & Adolescent Psychiatry*, 44(9), 789-790.

Woywodt A, Kiss A. Geophagia: the history of earth-eating (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1279487/>). *J R Soc Med.* 2002;95(3):143-146.