

Unlocking the potential of tio2 nanoparticles as anti-cancerous agent

Shreyoshi Pal¹, Pritha Pal^{2*}

¹M.Sc. student, Department of Biotechnology, School of Life Sciences, Swami Vivekananda University, Barrackpore, West Bengal- 700121

²Assistant Professor, Department of Microbiology, School of Life Sciences, Swami Vivekananda University, Barrackpore, West Bengal- 700121

*Corresponding author at: Department of Microbiology, School of Life Sciences, Swami Vivekananda University, Barrackpore, West Bengal- 700121. Corresponding Email: prithap@svu.ac.in

ABSTRACT

Titanium dioxide (TiO₂) nanoparticles have surfaced as a promising instrument in the realm of oncology, largely because of their distinct physicochemical attributes, biocompatibility and their capacity to generate reactive oxygen species (ROS) under certain conditions. Recent breakthroughs in nanotechnology have unveiled the potential of TiO₂ nanoparticles as effective anti-cancer agents: they facilitate targeted delivery, controlled drug release and increased cytotoxicity directed at cancer cells, while minimizing damage to healthy tissues. This review delves into the mechanisms that underlie the anti-cancer effects of TiO₂ nanoparticles, encompassing photocatalysis, the induction of oxidative stress and their role in synergizing with traditional therapies. The anti-cancer mechanisms of TiO₂ nanoparticles (which are composed of titanium dioxide) primarily involve the generation of reactive oxygen species (ROS) upon exposure to ultraviolet or visible light. This process induces oxidative stress, DNA damage and apoptosis in cancer cells. However, TiO₂ nanoparticles possess the potential to disrupt mitochondrial function and modulate signaling pathways that are crucial for cancer progression. Surface modifications and functionalization have, therefore, significantly enhanced their selectivity and efficacy; this

allows for targeted delivery while minimizing off-target effects. Although TiO₂ nanoparticles are acknowledged for their intrinsic anti-cancer properties, they offer significant potential across various applications (such as photodynamic therapy (PDT), drug delivery and bioimaging). When used in conjunction with chemotherapeutic agents, they demonstrate a synergistic effect, enhancing drug efficacy while concurrently reducing the necessary dosages. Recent advancements in nanotechnology have also facilitated their use as diagnostic tools, offering high-resolution imaging capabilities for early tumor detection. This versatility positions TiO₂ nanoparticles as a valuable asset in the ongoing fight against cancer; however, further research is essential to fully leverage their capabilities.

Keywords: *nanoparticles, biocompatibility, reactive oxygen species (ROS) , cytotoxicity , photocatalysis, oxidative stress , apoptosis , photodynamic therapy (PDT) , drug delivery and bioimaging.*

INTRODUCTION:

Cancer stands as one of the most widespread afflictions globally, impacting millions of individuals across diverse forms. In light of the considerable progress achieved in medical research, the quest for a definitive cure or universally effective therapy for cancer continues to be a challenging endeavor. Recent advancements in nanotechnology have unveiled intriguing possibilities for the improvement of cancer treatment and diagnosis. Nanoparticles, characterized as minuscule entities measuring less than 100 nanometers, have garnered considerable interest in the realms of health sciences and medicine owing to their distinctive attributes. Their diminutive dimensions, extensive surface area, and capacity for interaction at both cellular and molecular levels render them exceptionally suitable for precise therapeutic applications. It is essential to recognize that certain nanoparticles could potentially exert negative impacts on various animal cells, thereby underscoring the need for meticulous design and comprehensive testing within clinical environments.

Scholars are employing nanoparticles in diverse facets of cancer treatment to enhance efficacy and precision. In photodynamic therapy (PDT) and sonodynamic therapy (SDT), nanoparticles function as sensitizing agents, augmenting the production of reactive oxygen species to selectively eradicate cancer cells. In the realm of computed tomography (CT) imaging and radiation therapy, nanoparticles function as enhancement agents, augmenting the contrast and precision of imaging or amplifying the radiation dose administered to the tumor, all while mitigating harm to adjacent healthy tissues. Furthermore, there is ongoing development of nanoparticles for dual-mode imaging and therapy, wherein they function concurrently as contrast agents and therapeutic instruments. Titanium dioxide nanoparticles (TiO₂ NPs) stand out among the various types of nanoparticles due to their significant application in the medical field, particularly in cancer research. Titanium dioxide nanoparticles are employed in photodynamic therapy, sonodynamic therapy, and drug delivery systems owing to their compatibility with biological systems, photoreactive properties, and potential for functionalization aimed at the precise targeting of cancerous cells. The ongoing challenge of cancer as a global health issue underscores the imperative for the advancement of innovative therapeutics and treatment modalities. The advancements in nanotechnology, poised to transform cancer treatment, are set to be integral to the future landscape of oncology. Scholars are committed to enhancing nanoparticle-based methodologies to optimize their effectiveness while reducing potential hazards, with the ultimate goal of delivering more efficient and tailored cancer therapies. Nanotechnology, first introduced by Richard Feynman in 1959, stands as one of the most swiftly advancing fields of scientific exploration and technological progress worldwide. The utilization and deployment of nanoparticles are significantly oriented towards the domains of chemistry, energy, healthcare, and cosmetics (Santos et al., 2015). Nanoparticles can be composed of organic and inorganic components, including ferritins, liposomes, micelles, dendrimers, and nanoparticles endowed with magnetic properties, metals, and semiconductor nanoparticles, such as

oxides, nitrides, and sulfides, respectively (Nabi et al., 2019). Titania, an oxide of titanium, is classified as a nanoparticle within the realm of metal oxides, showcasing unique optical, thermal, electrical, and magnetic properties. Under standard conditions, TiO_2 remains insoluble in water and exhibits exceptional stability, characterized by a notable refractive index of $n = 2.4$, which makes it an ideal candidate for application as a white pigment. Titanium dioxide is generally found in three distinct crystalline polymorphs: anatase, rutile, and brookite. Among these three polymorphic forms, anatase and rutile display comparable properties; nonetheless, rutile possesses a lower density and demonstrates enhanced resistance to corrosion. Furthermore, brookite is infrequently observed owing to its instability at high temperatures, resulting in its conversion into rutile (Vijayalakshmi and Rajendran, 2012). The notable applications of nano TiO_2 encompass photocatalytic degradation and splitting, photovoltaic cells, electrochromic and electronic devices, along with sensing instruments. These applications have generated considerable interest and prompted extensive research and development efforts directed towards the synthesis of TiO_2 nanoparticles (Irshad et al., 2020, Mollavali et al., 2018, Narkevica et al., 2017, Pant et al., 2019).

Properties of TiO_2 nanoparticles:

Titanium dioxide nanoparticles (TiO_2 NPs) possess several distinctive properties that render them invaluable in diverse medical applications, particularly in cancer therapy. The following delineates the principal properties of TiO_2 NPs:

- 1. Photoreactivity:** TiO_2 NPs demonstrate robust photocatalytic activity, particularly under ultraviolet (UV) light. This characteristic allows for the effective generation of reactive oxygen species (ROS) upon exposure to light, a mechanism leveraged in photodynamic therapy (PDT) to eradicate cancer cells.
- 2. Biocompatibility:** TiO_2 NPs are generally regarded as biocompatible, indicating their ability to interact with biological systems without eliciting significant adverse effects. This trait is essential for their application in medical

fields, encompassing drug delivery and cancer treatment.

3. High Surface Area: Owing to their nanoscale dimensions, TiO₂ NPs feature a high surface area-to-volume ratio. This promotes efficient engagement with biological molecules and cells, thereby augmenting their efficacy in medical applications.

4. Chemical Stability: TiO₂ NPs exhibit chemical stability and resilience against oxidation and corrosion. This trait assures the retention of their functionality over time, which is vital for enduring medical applications.

5. Surface Functionalization: The surfaces of TiO₂ NPs can be readily modified with various chemical entities or biomolecules, facilitating targeted delivery to specific cells or tissues, including cancer cells.

6. Antimicrobial Properties: TiO₂ NPs have been demonstrated to possess antimicrobial characteristics, assisting in the prevention of infections when employed in medical devices or implants.

7. Optical Properties: TiO₂ NPs exhibit commendable optical attributes, including a high refractive index and substantial UV absorption. These characteristics enhance imaging modalities such as computed tomography (CT) scans.


8. Non-Toxicity: Given appropriate dosages and controlled conditions, TiO₂ NPs are relatively non-toxic, rendering them suitable for human applications. Nevertheless, their toxicity may vary based on size, shape, and surface enhancements.

These attributes establish TiO₂ NPs as versatile instruments in cancer therapy, notably in amplifying the effectiveness of existing treatments like PDT and serving as vehicles in drug delivery systems.


Mechanisms of Anti-Cancer Action :

Cancer cells demonstrate an increased vulnerability to oxidative stress in contrast to normal cells, largely as a result of their modified metabolic pathways and the heightened

production of reactive oxygen species (ROS). This attribute makes them appropriate candidates for therapeutic strategies aimed at further augmenting ROS levels.



Titanium dioxide (TiO₂) nanoparticles demonstrate significant photocatalytic activity upon exposure to light, particularly ultraviolet (UV) or near-UV light, which results in the generation of various ROS, including hydroxyl radicals, superoxide anions, and hydrogen peroxide. These reactive species possess the ability to damage essential cellular components such as DNA, proteins, and lipids, thereby initiating apoptosis, or programmed cell death, in cancerous cells.



In the context of cancer treatment, TiO₂ nanoparticles are utilized in two forms: pure and functionalized. Pure TiO₂ nanoparticles rely on their inherent photocatalytic capabilities

to produce ROS, while functionalized TiO_2 nanoparticles are enhanced with specific molecules or coatings to improve targeting precision and ROS production efficiency. This modification enhances their selectivity towards cancer cells, thereby minimizing potential off-target effects on healthy cells.

In the context of photodynamic therapy (PDT), the interplay between light activation and TiO_2 nanoparticles enables a precise and localized increase in reactive oxygen species (ROS) levels, thereby effectively suppressing tumor proliferation. This approach minimizes damage to healthy tissues, as the light activation can be accurately directed at the tumor site, thereby enhancing the production of reactive oxygen species in the vicinity of the nanoparticles.

In cellular level, the role of titanium dioxide nanoparticles (TiO_2 NPs) on cellular metabolism and survival, emphasizing the role of reactive oxygen species (ROS) at different doses of TiO_2 NPs.

Mechanism of Action of Low Concentrations of Titanium Dioxide Nanoparticles (TiO_2 NPs)

1. Moderate Reactive Oxygen Species (ROS)

Generation: At low concentrations, TiO_2 NPs generate a moderate quantity of reactive oxygen species (ROS) when subjected to light exposure. This induces a state of **mild oxidative stress** within cellular environments.

2. Stimulation of Antioxidant Defense Systems: The slight elevation of ROS levels activates the cell's **antioxidant defense mechanisms**. Crucial enzymes such as **superoxide dismutase (SOD)** are upregulated to neutralize ROS and avert cellular damage. SOD catalyzes the transformation of superoxide radicals into oxygen and hydrogen peroxide, which is less detrimental to cellular health.

3. Optimized Mitochondrial Function: In reaction to moderate oxidative stress, mitochondria adapt by enhancing their metabolic functions, specifically the **tricarboxylic acid (TCA) cycle**. This adaptation sustains cellular energy levels by ensuring a consistent supply of **adenosine triphosphate**

(ATP), the primary energy currency of the cell.

4. Cell Viability: The cell successfully maintains its metabolic equilibrium and energy production, enabling it to endure moderate oxidative stress. This resilience is conducive to continuing normal cellular function or facilitating recovery from mild stress conditions.

Mechanism of Action of High Concentrations of Titanium Dioxide Nanoparticles (TiO₂ NPs)

1. The elevated production of reactive oxygen species (ROS) is observed with high concentrations of TiO₂ nanoparticles, leading to a marked increase in ROS levels. This escalation results in significant oxidative stress, exceeding the cell's natural antioxidant defenses.

Mitochondrial Impairment: The significant elevation in reactive oxygen species (ROS) causes harm to the mitochondria, leading to a disturbance in their membrane potential (MMP) and ultimately resulting in mitochondrial dysfunction. This dysfunction impedes the mitochondria's capacity to efficiently execute the TCA cycle, resulting in a significant decline in ATP production.

3. The release of cytochrome C occurs as a consequence of mitochondrial damage, leading to its presence in the cytoplasm. Cytochrome C is integral to the electron transport chain, and its release marks the commencement of mitochondrial-induced apoptosis, also known as the intrinsic apoptotic pathway.

4. The induction of the apoptotic pathway occurs when cytochrome C is released, leading to the activation of caspase 9, which in turn triggers caspase 3. Caspases serve as essential mediators within the apoptotic process. Caspase 9 serves as the initiator of the apoptotic signaling cascade, whereas caspase 3 carries out the cell death program through the cleavage of various cellular substrates.

5. The activation of the caspase cascade leads to apoptosis, a sophisticated mechanism of programmed cell death. Apoptosis is defined by a reduction in cell size, fragmentation of DNA, and the subsequent uptake by adjacent cells or phagocytes, which functions to mitigate inflammation. It can be inferred that:

1. Low concentrations of TiO_2 NPs augment the cell's ability to cope with oxidative stress, consequently improving cellular survival and metabolic stability.
2. Elevated levels of TiO_2 NPs, on the other hand, induce oxidative stress that surpasses the cellular capacity for management, leading to mitochondrial impairment, the initiation of apoptotic pathways, and ultimately, cell demise. The dual mechanism of action of TiO_2 NPs positions them as a significant contributor to cancer therapy, aiming to induce apoptosis in malignant cells while safeguarding the integrity of normal cells.

Potential Applications in Cancer Therapy: Titanium dioxide nanoparticles (NPs) find application across a wide spectrum of fields in medicine and life sciences. These nanoparticles act as vehicles for pharmaceuticals, especially anticancer agents, and operate as facilitators for photodynamic therapy (PDT) or photothermal therapy (PTT) in the management of solid tumors. Titanium dioxide nanoparticles have been adeptly utilized as carriers for a range of anticancer therapeutics, such as paclitaxel, doxorubicin, daunorubicin, temozolomide, and camptothecin. TiO_2 nanoparticles can be used to *directly target and kill cancer cells when activated by light*. This approach is particularly useful for *superficial cancers* (e.g., *skin cancer*) where light penetration is sufficient. Titanium dioxide nanoparticles (TiO_2 NPs) exhibit a diverse array of potential applications in cancer therapy, attributable to their unique properties.

Principal applications encompass:

1. Photodynamic Therapy (PDT): Titanium dioxide nanoparticles generate reactive oxygen species (ROS) when subjected to light, resulting in oxidative harm to malignant cells while safeguarding surrounding healthy tissue.
2. Sonodynamic Therapy (SDT): Analogous to Photodynamic Therapy (PDT), this methodology employs ultrasound to stimulate TiO_2 nanoparticles, thereby producing reactive oxygen species (ROS) to proficiently address deep-seated tumors.

3. **Drug Delivery Systems:** TiO_2 NPs enable the precise administration of anticancer pharmaceuticals to tumor locations, thereby improving therapeutic effectiveness and minimizing undesirable side effects.

4. **The enhancement of radiation therapy** through TiO_2 nanoparticles is noteworthy, as they significantly increase the production of reactive oxygen species during treatment, thereby augmenting its efficacy in targeting cancer cells.

5. **CT Imaging:** The incorporation of TiO_2 nanoparticles significantly improves tumor visibility in CT scans, thereby facilitating precise diagnosis and enabling targeted therapeutic interventions.

6. **Dual-Mode Imaging and Therapy:** TiO_2 NPs serve a dual purpose as both imaging agents and therapeutic tools, facilitating the seamless integration of diagnosis and treatment.

7. **Induction of Apoptosis:** TiO_2 NPs generate oxidative stress within cancer cells, thereby initiating programmed cell death (apoptosis) and inhibiting tumor proliferation, all while preserving the integrity of normal cells.

LIMITATION AND CHALLENGES:

While the potential applications of titanium dioxide nanoparticles (TiO_2 NPs) in cancer therapy are indeed promising, there are several noteworthy challenges that must be confronted:

1. **Toxicity:** Although TiO_2 NPs are typically regarded as biocompatible, there exists the potential for toxicity when administered at high doses or subjected to extended exposure.

2. **Targeting Efficiency:** The precise identification of cancer cells while safeguarding healthy tissues continues to present a significant challenge.

3. **Regulatory Issues:** The journey toward clinical application for TiO_2 NPs is fraught with rigorous approval processes, which elicit apprehensions regarding safety and environmental ramifications.

4. Aggregation: Within biological contexts, the aggregation of TiO_2 nanoparticles may occur, potentially reducing their efficacy and obstructing their capacity to traverse barriers, including the blood-brain barrier.

5. Light Penetration: The restricted capacity of light to infiltrate during photodynamic therapy diminishes the effectiveness of treatment for tumors situated deeper within tissues.

6. Scalability: The endeavor of producing these nanoparticles in substantial quantities while ensuring uniform properties poses a considerable challenge.

7. Absence of Standardization: The variability in nanoparticle properties presents significant challenges for their integration into clinical applications.

8. Immune Response: The aggregation of TiO_2 nanoparticles has the potential to elicit immune responses, consequently influencing their safety and therapeutic effectiveness.

Future prospect: TiO_2 NPs are highly photoactive catalytic agent, with their affordable price, stability, it can be a hope for future cancer therapy. Despite having several limitations, these limitations underscore the necessity for further research to optimize TiO_2 NPs for cancer treatment.

CONCLUSION:

In summary, TiO_2 nanoparticles exhibit significant promise as anti-cancer agents, attributed to their photocatalytic properties, capacity to produce reactive oxygen species, and compatibility with biological systems. Despite the ongoing challenges related to safe delivery and precise activation, the progress in nanoparticle engineering and medical technology is poised to broaden their utilization in cancer treatment. Subsequent investigations will concentrate on overcoming existing constraints, examining the potential of combination therapies, and progressing towards clinical trials to validate TiO_2 nanoparticles as a credible alternative in the field of oncology.

REFERENCES:

- Çeşmeli, S., & Biray Avci, C. (2019). Application of titanium dioxide (TiO₂) nanoparticles in cancer therapies. *Journal of drug targeting*, 27(7), 762-766.
- Raja, G., Cao, S., Kim, D. H., & Kim, T. J. (2020). Mechanoregulation of titanium dioxide nanoparticles in cancer therapy. *Materials Science and Engineering: C*, 107, 110303.
- Robertson TA, Sanchez WY, Roberts MS: Are commercially available nanoparticles safe when applied to the skin? *J Biomed Nanotechnol* 2010, 6: 452-468. 10.1166/jbn.2010.1145
- Nasikhudin^{1,2}, M Diantoro², A Kusumaatmaja¹, K Triyana¹, Study on Photocatalytic Properties of TiO₂ Nanoparticle in various pH condition The International Conference on Theoretical and Applied Physics IOP Publishing IOP Conf. Series: Journal of Physics: Conf. Series 1011 (2018) 012069
- Ziental, D., Czarczynska-Goslinska, B., Mlynarczyk, D. T., Glowacka-Sobotta, A., Stanisz, B., Goslinski, T., & Sobotta, L. (2020). Titanium dioxide nanoparticles: prospects and applications in medicine. *Nanomaterials*, 10(2), 387.
- Zarzzeka, C., Goldoni, J., Marafon, F., Sganzerla, W. G., Forster-Carneiro, T., Bagatini, M. D., & Colpini, L. M. S. (2023). Use of titanium dioxide nanoparticles for cancer treatment: A comprehensive review and bibliometric analysis. *Biocatalysis and Agricultural Biotechnology*, 50, 102710.
- Fujiwara, R., Luo, Y., Sasaki, T., Fujii, K., Ohmori, H., & Kuniyasu, H. (2015). Cancer therapeutic effects of titanium dioxide nanoparticles are associated with oxidative stress and cytokine induction. *Pathobiology*, 82(6), 243-251.
- Sargazi, S., Simge, E. R., Gelen, S. S., Rahdar, A., Bilal, M., Arshad, R., ... & Pandey, S. (2022). Application of titanium dioxide nanoparticles in photothermal and photodynamic therapy of cancer: An updated and comprehensive review. *Journal of drug delivery science and technology*, 75, 103605.
- Zarzzeka, C., Goldoni, J., Marafon, F., Sganzerla, W. G., Forster-Carneiro, T., Bagatini, M. D., & Colpini, L. M. S. (2023). Use of titanium dioxide nanoparticles for cancer treatment: A comprehensive review and bibliometric analysis. *Biocatalysis and Agricultural Biotechnology*, 50, 102710.

Smith, L., Kuncic, Z., Ostrikov, K., & Kumar, S. (2012). Nanoparticles in cancer imaging and therapy. *Journal of Nanomaterials*, 2012(1), 891318.

Kim, S., Im, S., Park, E. Y., Lee, J., Kim, C., Kim, T. I., & Kim, W. J. (2020). Drug-loaded titanium dioxide nanoparticle coated with tumor targeting polymer as a sonodynamic chemotherapeutic agent for anti-cancer therapy. *Nanomedicine: Nanotechnology, Biology and Medicine*, 24, 102110.