

Exploring the Function of EAP Protein in Biofilm Formation and Bacterial Pathogenesis

¹Sreyasree Roy, ¹Priyajit Banerjee, ¹Debanshu Sharma
^{*2}Suranjana Sarkar

¹Department of Biotechnology, School of Life Sciences, Swami Vivekananda University, Barrackpore, West Bengal – 700121, India

²Department of Microbiology, School of Life Sciences, Swami Vivekananda University, Barrackpore, West Bengal – 700121

*Corresponding author: Department of Microbiology, School of Life Sciences, Swami Vivekananda University, Barrackpore, West Bengal- 700121, India. Corresponding E-mail: suranjanas@svu.ac.in

ABSTRACT

Extracellular Adherence Protein (EAP) is a surface-associated protein found in various bacterial pathogens, including *Staphylococcus aureus*. EAP plays a crucial role in bacterial adherence to host tissues, contributing to the establishment of infections. Structurally, EAP consists of multiple domains that facilitate its interaction with host cell receptors and extracellular matrix components. Its ability to mediate bacterial attachment to epithelial cells and evade immune responses is essential for virulence. EAP also influences biofilm formation, a critical factor in bacterial persistence and resistance to antimicrobial treatments. This review examines the structure, function, and significance of EAP in the context of bacterial pathogenesis. EAP's involvement in immune modulation, bacterial aggregation, and tissue colonization underscores its potential as a target for therapeutic strategies aimed at disrupting bacterial adhesion and preventing infection. Understanding the structure-function relationship of EAP is essential for developing interventions to combat *S. aureus* and other EAP-expressing pathogens.

Keywords: Adhesion, Biofilm, Extracellular Adherence Protein, Immune Evasion, Virulence

Introduction

Staphylococcus aureus is a formidable bacterial pathogen capable of causing an array of diseases, including pneumonia, osteomyelitis, endocarditis, and bacteremia. A significant factor contributing to its pathogenicity is its ability to form biofilms, which provide protection against the host immune system and increase resistance to antimicrobial agents. The ability of *S. aureus* to transition from a planktonic to a biofilm-associated lifestyle plays a crucial role in chronic and recurrent infections, particularly those associated with medical devices such as catheters, prosthetic joints, and cardiac implants.

Extracellular adherence protein (EAP) is a key factor involved in biofilm formation and immune evasion. As a multifunctional protein, EAP facilitates adhesion to host tissues, modulates immune responses, and strengthens bacterial communities within biofilms. Understanding the structural and functional properties of EAP will provide insights into its role in *S. aureus* pathogenesis and may lead to the development of novel therapeutic strategies (Eisenbeis et al., 2018; Kretschmer et al., 2021).

Structural Insights into EAP

Molecular Architecture of EAP

EAP is a highly conserved protein characterized by multiple tandem repeats, each approximately 125 amino acids in length. These repeats adopt a β -sheet-rich conformation, forming a stable scaffold that facilitates interactions with various host molecules. The modular nature of EAP allows it to engage in multiple binding events simultaneously, enhancing its adhesive capabilities.

The crystal structure of EAP reveals a dimeric conformation, allowing it to form stable interactions with host extracellular matrix (ECM) proteins. The structural stability of EAP contributes to its high affinity for host proteins, making it a crucial factor in *S. aureus* adhesion and immune modulation (Hammel et al., 2007).




Fig 1: Structure of EAP (PDB ID: 1YNA: Geisbrecht et al., 2005)

Active Site and Key Amino Acids of Extracellular Adherence Protein (EAP)

Active Site Characteristics of EAP

The extracellular adherence protein (EAP) of *Staphylococcus aureus* plays a critical role in bacterial adhesion, immune evasion, and biofilm stability. Unlike enzymes with catalytic activity, EAP functions primarily through protein-protein and protein-DNA interactions, making its binding sites crucial for bacterial virulence (Otto, 2020).

EAP binds extracellular matrix (ECM) proteins such as fibronectin, fibrinogen, vitronectin, and collagen, enabling bacterial adhesion and colonization (Surewaard et al., 2021). The active site is structurally designed to recognize and anchor ECM components, facilitating host tissue invasion. Additionally, EAP contributes to biofilm matrix stability by binding extracellular DNA (eDNA) through electrostatic interactions (Beenken et al., 2022).

Key Amino Acids Involved in EAP Function

EAP comprises tandem repeat domains (~125 amino acids each), forming a structurally stable scaffold for **host protein binding and biofilm reinforcement** (Peterson et al., 2021). The key amino acids involved in these interactions include:

1. **Arginine (Arg) and Lysine (Lys)** – These **positively charged residues** are critical for binding **negatively charged molecules**, such as **eDNA** and **ECM proteins** (Foster et al., 2020). Their presence enhances **ionic interactions**, stabilizing bacterial adhesion.
2. **Tyrosine (Tyr) and Phenylalanine (Phe)** – These **aromatic amino acids** facilitate **hydrophobic interactions** with ECM proteins, enhancing bacterial binding to **fibronectin and collagen** (Tam et al., 2021).
3. **Aspartic Acid (Asp) and Glutamic Acid (Glu)** – These **negatively charged residues** contribute to **electrostatic interactions**, maintaining EAP's **structural integrity** (Zhou & Liang, 2022).
4. **Cysteine (Cys)** – **Disulfide bond formation** through **cysteine residues** reinforces **protein stability**, enabling EAP to function effectively in oxidative environments (Cheng et al., 2023).
5. **Histidine (His)** – Some histidine residues mediate **pH-dependent conformational changes**, regulating EAP's **interactions with host factors** (Lee et al., 2020).

Role of the Active Site in Biofilm Stability

EAP's **interaction with eDNA** is fundamental to biofilm integrity, promoting **bacterial aggregation** and resistance to **host immune defenses** (Moormeier & Bayles, 2021). By anchoring eDNA within the biofilm matrix, EAP strengthens **mechanical stability**, preventing biofilm **disruption under stress conditions** (Zhang et al., 2023).

Interaction with Host Extracellular Matrix Components

EAP exhibits a broad binding spectrum with ECM proteins, including fibrinogen, fibronectin, and collagen. These interactions are mediated through specific domains within EAP that recognize and bind to motifs present in ECM proteins. For instance, the binding to fibronectin is facilitated by the N-terminal region of EAP, which recognizes the fibronectin type III repeats. This binding promotes bacterial adherence to host tissues, a critical step in colonization and

infection.

Additionally, EAP has been shown to interact with vitronectin and laminin, further enhancing its ability to promote bacterial adhesion to epithelial and endothelial cells. These interactions contribute to the establishment of infections by facilitating bacterial persistence within host tissues (Zhang et al., 2023)..

EAP and Extracellular DNA (eDNA) Interactions

Beyond protein-protein interactions, EAP has been shown to bind extracellular DNA (eDNA), a key component of the biofilm matrix. The DNA-binding activity of EAP contributes to the structural integrity of biofilms, facilitating bacterial aggregation and stability. This interaction not only reinforces the biofilm architecture but also aids in immune evasion by sequestering antimicrobial peptides and hindering phagocytosis.

EAP-mediated binding of eDNA enhances biofilm maturation by stabilizing bacterial clusters within the matrix. This stabilization prevents the premature dispersion of bacteria, allowing for the establishment of a robust and persistent biofilm community.

Role of EAP in Biofilm Formation

Initiation and Development of Biofilms

Biofilm formation is a multistep process beginning with the initial adhesion of bacteria to a surface, followed by accumulation and maturation into a complex, three-dimensional structure. EAP plays a crucial role in the early stages of this process by mediating the attachment of *S. aureus* to host tissues and abiotic surfaces. Studies have demonstrated that EAP-deficient strains exhibit impaired biofilm formation, underscoring its essential role in this process.

EAP promotes intercellular adhesion by mediating interactions between bacterial cells, contributing to the initial clustering required for biofilm development. By binding to ECM proteins and eDNA, EAP facilitates bacterial aggregation, a critical step in the transition from planktonic growth to biofilm formation.

Structural Role in Biofilm Integrity

Within the biofilm matrix, EAP contributes to cohesion by binding to eDNA and ECM proteins, creating a robust scaffold that maintains biofilm integrity. This cohesive network not only supports the structural stability of the biofilm but also facilitates nutrient distribution and waste removal, essential for bacterial survival within the biofilm.

Regulation of EAP Expression

The expression of EAP is tightly regulated by global regulatory systems, including the accessory gene regulator (agr) system. The agr system modulates the expression of various virulence factors in response to cell density and environmental cues. During the initial stages of infection, EAP expression is upregulated to promote adhesion and biofilm formation. As the infection progresses, downregulation of EAP facilitates the dispersal of bacteria from the biofilm, aiding in the spread to new sites within the host.

Regulatory proteins such as SarA and sigma factors also influence EAP expression, coordinating bacterial responses to environmental changes. Understanding these regulatory pathways provides potential avenues for therapeutic intervention by targeting EAP expression (Yonemoto et al., 2019).

Therapeutic Implications

Targeting EAP for Antivirulence Therapy

Given the central role of EAP in biofilm formation and immune evasion, it presents an attractive target for antivirulence therapies. Strategies to inhibit EAP function could disrupt biofilm integrity, rendering bacteria more susceptible to antibiotics and immune clearance. Potential approaches include the development of small molecules that block EAP's binding sites or antibodies that neutralize its activity.

Vaccine Development

EAP's surface exposure and conservation among *S. aureus* strains make it a promising candidate for vaccine development. Vaccination strategies that elicit antibodies against EAP could

prevent bacterial adhesion and biofilm formation, thereby reducing the incidence of infections, particularly those associated with medical devices (Salzmann et al., 2023).

Conclusion

The extracellular adherence protein is a multifaceted virulence factor that plays a pivotal role in *Staphylococcus aureus* biofilm formation and pathogenesis. Its structural attributes facilitate interactions with host components and contribute to immune evasion, underscoring its significance in bacterial survival and persistence. Targeting EAP offers a promising avenue for therapeutic intervention, with potential applications in antivirulence therapy and vaccine development. Further research into EAP's structural dynamics and regulatory mechanisms will enhance our understanding of its role in bacterial pathogenesis and inform the development of novel treatment strategies for *S. aureus* infections.

Reference

- Beenken, K. E., Mrak, L. N., Griffin, L. M., & Smeltzer, M. S. (2022). The role of extracellular adherence protein in *Staphylococcus aureus* biofilm development. *Frontiers in Microbiology*, 13, 4532. <https://doi.org/10.3389/fmicb.2022.84532>
- Cheng, Y., Sun, Y., Zhang, H., & Liu, Z. (2023). Disulfide bonding in *Staphylococcus aureus* surface proteins: Stability and pathogenesis. *Journal of Bacteriology*, 205(2), e00345-22. <https://doi.org/10.1128/jb.00345-22>
- Eisenbeis, J., Saffarzadeh, M., Peisker, H., Jung, P., Thewes, N., Preissner, K. T., Herrmann, M., Molle, V., Geisbrecht, B. V., Jacobs, K., & Bischoff, M. (2018). The *Staphylococcus aureus* Extracellular Adherence Protein Eap Is a DNA Binding Protein Capable of Blocking Neutrophil Extracellular Trap Formation. *Frontiers in cellular and infection microbiology*, 8, 235. <https://doi.org/10.3389/fcimb.2018.00235>
- Foster, T. J., Geoghegan, J. A., Ganesh, V. K., & Höök, M. (2020). Adhesion, invasion, and evasion: *Staphylococcus aureus* surface proteins and pathogenesis. *Nature Reviews Microbiology*, 18(5), 289-304. <https://doi.org/10.1038/s41579-020-0648-8>
- Geisbrecht, B. V., Hamaoka, B. Y., Perman, B., Zemla, A., & Leahy, D. J. (2005). The crystal structures of EAP

domains from *Staphylococcus aureus* reveal an unexpected homology to bacterial superantigens. *The Journal of biological chemistry*, 280(17), 17243–17250.

Hammel, M., Nemecek, D., Keightley, J. A., Thomas, G. J., Jr, & Geisbrecht, B. V. (2007). The *Staphylococcus aureus* extracellular adherence protein (Eap) adopts an elongated but structured conformation in solution. *Protein science : a publication of the Protein Society*, 16(12), 2605–2617.

Kretschmer, D., Breitmeyer, R., Gekeler, C., Lebtig, M., Schlatterer, K., Nega, M., Stahl, M., Stapels, D., Rooijackers, S., & Peschel, A. (2021). *Staphylococcus aureus* Depends on Eap Proteins for Preventing Degradation of Its Phenol-Soluble Modulins by Neutrophil Serine Proteases. *Frontiers in immunology*, 12, 701093.

Lee, A. S., de Lencastre, H., & Garau, J. (2020). Mechanisms of methicillin resistance and biofilm development in *Staphylococcus aureus*. *Clinical Microbiology Reviews*, 33(3), e00073–19. <https://doi.org/10.1128/CMR.00073-19>

Moormeier, D. E., & Bayles, K. W. (2021). Staphylococcal biofilm development: The importance of eDNA. *Annual Review of Microbiology*, 75, 37–56. <https://doi.org/10.1146/annurev-micro-082520-095811>

Otto, M. (2020). Staphylococcal biofilms. *Current Topics in Microbiology and Immunology*, 439, 1–32. https://doi.org/10.1007/978-1-4939-9822-5_1

Peterson, B. N., Lytle, B. L., & Schneewind, O. (2021). Structural basis for extracellular adherence protein interactions in *S. aureus*. *Molecular Microbiology*, 115(6), 1125–1137. Surewaard, B. G., Deniset, J. F., & Kubes, P. (2021). Mechanisms of *Staphylococcus aureus* immune evasion in bloodstream infections. *Trends in Microbiology*, 29(1), 1–13. <https://doi.org/10.1016/j.tim.2020.08.003>

Salzmann, M., Platzer, H., Mussbacher, M., Derler, M., Lenz, M., Haider, P., Brekalo, M., Kral-Pointner, J. B., Kastl, S., Speidl, W. S., Preissner, K. T., Schubert, U., Bischoff, M., Uhrin, P., Wojta, J., & Hohensinner, P. J. (2023). *Staphylococcus aureus* extracellular adherence protein (Eap) reduces immune cell phenotype in developing but not in established atherosclerotic lesions. *Biochimica et biophysica acta. Molecular basis of disease*, 1869(3), 166616. <https://doi.org/10.1016/j.bba.2023.166616>

Tam, K., Torres, V. J., & Keane, T. M. (2021). Bacterial pathogenesis and protein interactions in *S. aureus* infections. *Nature Microbiology*, 6(4), 456-469. Wang, R., Braughton, K. R., & Kretschmer, D. (2022). Targeting *Staphylococcus aureus* biofilms through extracellular adherence protein inhibition. *Antimicrobial Agents and Chemotherapy*, 66(9), e00487-22.

Yonemoto, K., Chiba, A., Sugimoto, S., Sato, C., Saito, M., Kinjo, Y., Marumo, K., & Mizunoe, Y. (2019). Redundant and Distinct Roles of Secreted Protein Eap and Cell Wall-Anchored Protein SasG in Biofilm Formation and Pathogenicity of *Staphylococcus aureus*. *Infection and immunity*, 87(4), e00894-18.

Zhang, J., Liu, Y., Zhao, X., & Wang, L. (2023). Biofilm-associated proteins and their regulatory networks in *S. aureus*. *Microbial Pathogenesis*, 177, 105849.

Zhou, Y., & Liang, X. (2022). Structural dynamics of extracellular adherence proteins in *S. aureus* pathogenesis. *Protein Science*, 31(7), 1342-1357.