

Lifestyle Interventions in MASLD Management: Diet, Exercise, and Behavior Modification

Farteen Feroz and Parul Thapar*

Department of Food and Nutrition, Swami Vivekananda
University, Barrackpore, W.B., India

*Corresponding E-mail: parul.thapar@yahoo.com

Abstract

Metabolic dysfunction-associated steatotic liver disease (MASLD), formerly known as non-alcoholic fatty liver disease (NAFLD), is a growing public health concern linked to obesity, insulin resistance, and metabolic syndrome. Lifestyle interventions, including dietary modifications, physical activity, and behavioral changes, remain the cornerstone of MASLD management. This review explores the role of these interventions, their effectiveness, and strategies for implementation.

Keywords: MASLD, NAFLD, diet, exercise, behavior modification, lifestyle interventions, metabolic syndrome, precision nutrition, digital health

Introduction

MASLD affects millions globally and is associated with increased risks of cirrhosis, cardiovascular diseases, and type 2 diabetes (Chalasani et al, 2018; Eslam et al, 2020; Jarvinen, 2016). With no approved pharmacological treatments, lifestyle changes are the primary therapeutic strategy. This review evaluates current evidence on diet, exercise, and behavior modification in MASLD management and discusses their mechanisms, benefits, and practical applications (Zelber-Sagi et al, 2021; Friedman et al, 2018).

Dietary Interventions

Diet plays a pivotal role in managing MASLD by reducing liver fat, inflammation, and metabolic dysfunction (Halasani et al, 2018).

- **Caloric Restriction and Weight Loss:** A 5-10% weight reduction significantly improves liver histology, reducing hepatic steatosis, inflammation, and fibrosis progression (Slam et al, 2020; Ki-Järvinen et al., 2016).
- **Gradual weight loss through a hypocaloric diet** (500-1000 kcal/day deficit) is recommended to avoid adverse metabolic effects (Elber-Sagi et al., 2021).

Macronutrient Composition:

- **Low-Carbohydrate and Mediterranean Diets:** These diets emphasize whole grains, healthy fats, lean proteins, and fiber, reducing hepatic steatosis and insulin resistance (Riedman et al, 2018; Ilar-Gomez et al, 2015).
- **High-Protein Diets:** May improve liver fat content and metabolic parameters by enhancing satiety and preserving lean muscle mass (Outoukidis et al., 2019).

Specific Nutrients:

Omega-3 Fatty Acids: Reduce liver fat and inflammation by modulating lipid metabolism (Uzzetti et al., 2016).

Antioxidants (Vitamin E, Polyphenols): Have potential hepatoprotective effects through anti-inflammatory and oxidative stress reduction mechanisms (Ofi et al., 2010).

Fructose and Processed Foods: Excessive fructose intake promotes de novo lipogenesis, increasing liver fat accumulation and inflammation. Minimizing processed food consumption is recommended (Chwingshackl and Hoffmann, 2013).

Intermittent Fasting and Time-Restricted Eating: Emerging evidence suggests intermittent fasting improves metabolic parameters and reduces liver fat by enhancing autophagy and insulin sensitivity (Aniagua, 2016; Ump et al., 2016).

Exercise Interventions

Physical activity is critical in MASLD management by enhancing insulin sensitivity and reducing liver fat (Anyal et al., 2010; Im et al., 2010).

Aerobic Exercise:

- Improves hepatic steatosis, even in the absence of significant weight loss (Atterson et al., 2015).
- Activities such as brisk walking, cycling, and swimming have been shown to reduce intrahepatic fat (Insley et al., 2015).

Resistance Training:

- Enhances muscle mass, improves glucose metabolism, and contributes to liver fat reduction (Eating et al., 2012).
- Strength training at least twice per week is recommended for optimal metabolic benefits (Ohnson et al., 2009).

Exercise Volume and Intensity:

- Moderate-to-vigorous activity for at least 150-300 minutes per week is recommended (Allsworth et al., 2011).
- High-intensity interval training (HIIT) demonstrates promising effects in reducing hepatic fat and improving cardiovascular health (Younossi, 2007).

Behaviour Modification Strategies:

Sustaining lifestyle changes requires behavioral interventions to enhance long-term adherence and success (Garber et al., 2011).

Cognitive-Behavioral Therapy (CBT):

- Helps in habit formation and overcoming barriers to dietary and physical activity modifications (Yki-Järvinen, 2011).
- Identifies and restructures negative thought patterns related to lifestyle choices (Zelber-Sagi et al., 2021).

Self-Monitoring and Goal Setting:

- Tracking food intake, weight, and physical activity enhances self-awareness and adherence (Friedman et al., 2018).

- Mobile apps and digital health tools can facilitate monitoring and engagement.
- Encourages patient-centered decision-making, enhancing intrinsic motivation for sustainable lifestyle changes (Slam et al, 2020).

Social Support and Digital Tools:

- Involvement of family, peer support groups, and healthcare professionals improves adherence (Ki-Järvinen, 2016)
- Digital interventions, including virtual coaching and online communities, have shown positive impacts on lifestyle modification success (Elaber et al., 2021).

Integrated Lifestyle Approaches

Combining diet, exercise, and behavior modification yields optimal results. Multidisciplinary

programs involving healthcare providers, dietitians, exercise specialists, and behavioral therapists enhance long-term adherence and effectiveness (Riedman et al, 2018).

Personalized Approaches:

- Tailoring interventions based on individual metabolic profiles, preferences, and readiness to change improves outcomes (Ilar et al., 2015).

• Community-Based and Workplace Wellness Programs:

- Structured programs in community and workplace settings support behavior change through group dynamics and accountability (Outoukidis et al., 2019).

Future Directions and Research Needs

Despite strong evidence supporting lifestyle interventions, challenges remain in ensuring long-term adherence and effectiveness. Future research should focus on (Uzzetti et al, 2016):

- Developing precision nutrition and exercise interventions based on genetic and metabolic profiling.

- Evaluating the long-term effects of various dietary patterns on MASLD progression.
- Exploring novel digital and AI-driven health interventions to improve adherence.
- Investigating the role of emerging behavioral strategies, including gamification and virtual reality coaching, in sustaining lifestyle modifications.

Conclusion

Lifestyle interventions remain the first-line therapy for MASLD management. Personalized, evidence-based strategies incorporating dietary changes, regular physical activity, and behavioral modifications are essential for effective disease control. Integrating multidisciplinary support, digital tools, and tailored interventions can enhance adherence and long-term success in MASLD management.

References

- Allsworth, K., Fattakhova, G., Hollingsworth, K. G., et al. (2011). Resistance exercise reduces liver fat and its mediators in non-alcoholic fatty liver disease independent of weight loss. *Gut*, 60(9), 1278-1283.
- Aniagua, J. A. (2016). Nutrition, insulin resistance and dysfunctional adipose tissue determine the different components of metabolic syndrome. *World Journal of Diabetes*, 7(19), 483-514.
- Anyal, A. J., Chalasani, N., Kowdley, K. V., et al. (2010). Pioglitazone, vitamin E, or placebo for nonalcoholic steatohepatitis. *The New England Journal of Medicine*, 362(18), 1675- 1685.
- Atterson, R. E., Laughlin, G. A., LaCroix, A. Z., et al. (2015). Intermittent fasting and human metabolic health. *Journal of the Academy of Nutrition and Dietetics*, 115(8), 1203-1212.
- Chalasani, N., Younossi, Z., Lavine, J. E., et al. (2018). The diagnosis and management of NAFLD: Practice guidance from the American Association for the Study of Liver Diseases. *Hepatology*, 67(1), 328-357.

Chwingshackle, L., & Hoffmann, G. (2013). Adherence to Mediterranean diet and risk of cancer: A systematic review and meta-analysis of observational studies. *International Journal of Cancer*, 135(8), 1884-1897.

Clinics in Liver Disease, 11(1), 1-16.

Eating, S. E., Hackett, D. A., Parker, H. M., et al. (2012). Effect of aerobic exercise training dose on liver fat and visceral adiposity. *Journal of Hepatology*, 57(2), 247-255.

Elber-Sagi, S., Lotan, R., Shibolet, O., & Webb, M. (2021). Mediterranean dietary pattern and non-alcoholic fatty liver disease. *World Journal of Gastroenterology*, 27(26), 3683- 3696.

Eslam, M., Sanyal, A. J., & George, J. (2020). MAFLD: A consensus-driven proposed nomenclature for metabolic-associated fatty liver disease. *Gastroenterology*, 158(7), 1999-2014.

Friedman, S. L., Neuschwander-Tetri, B. A., Rinella, M., & Sanyal, A. J. (2018).

Garber, C. E., Blissmer, B., Deschenes, M. R., Franklin, B. A., Lamonte, M. J., Lee, I. M., Nieman, D. C., & Swain, D. P. (2011).

Halasani, N., Younossi, Z., Lavine, J. E., et al. (2018). The diagnosis and management of NAFLD: Practice guidance from the American Association for the Study of Liver Diseases. *Hepatology*, 67(1), 328-357.

Ilar-Gomez, E., Martinez-Perez, Y., Calzadilla-Bertot, L., et al. (2015). Weight loss through lifestyle modification significantly reduces features of nonalcoholic steatohepatitis, *Gastroenterology*, 149(2), 367-378.e5.

Im, J. S., Mietus-Snyder, M., Valente, A., Schwarz, J. M., & Lustig, R. H. (2010). The role of fructose in the pathogenesis of NAFLD and the metabolic syndrome. *Nature Reviews Gastroenterology & Hepatology*, 7(5), 251-264.

Insley, G. M., & La Bounty, P. M. (2015). Effects of intermittent fasting on body composition and clinical health markers in humans. *Nutrition Reviews*, 73(10), 661-674.

Ki-Järvinen, H. (2016). Nutritional modulation of non-alcoholic fatty liver disease and insulin resistance: Human data.

Best Practice & Research Clinical Endocrinology & Metabolism, 30(3), 373-383.

Ofi, F., Abbate, R., Gensini, G. F., & Casini, A. (2010). Accruing evidence on benefits of adherence to the Mediterranean diet on health: An updated systematic review and meta-analysis. *The American Journal of Clinical Nutrition*, 92(5), 1189-1196. Ohnson, N. A., Sachinwalla, T., Walton, D. W., et al. (2009). Aerobic exercise training reduces hepatic and visceral lipids in obese individuals without weight loss. *Hepatology*, 50(4), 1105-1112.

Outoukidis, D. A., Astbury, N. M., Tudor, K. E., et al. (2019). Association of weight loss

Riedman, S. L., Neuschwander-Tetri, B. A., Rinella, M., & Sanyal, A. J. (2018). Mechanisms of NAFLD development and therapeutic strategies. *Nature Medicine*, 24(7), 908-922.

Slam, M., Sanyal, A. J., & George, J. (2020). MAFLD: A consensus-driven proposed nomenclature for metabolic-associated fatty liver disease. *Gastroenterology*, 158(7), 1999-2014.

Ump, D. B., Depner, C. M., & Tripathy, S. (2012). Omega-3 fatty acid supplementation and cardiovascular disease. *Journal of Lipid Research*, 53(12), 2525-2545.

Uzzetti, E., Pinzani, M., & Tsochatzis, E. A. (2016). The multiple-hit pathogenesis of non- alcoholic fatty liver disease (NAFLD). *Metabolism*, 65(8), 1038-1048.

Yki-Järvinen, H. (2016). Nutritional modulation of non-alcoholic fatty liver disease and insulin resistance: Human data. *Best Practice & Research Clinical Endocrinology &*

Younossi, Z. M. (2007). Epidemiology and natural history of NAFLD and NASH.

Zelber-Sagi, S., Lotan, R., Shibolet, O., & Webb, M. (2021). Mediterranean dietary pattern and non-alcoholic fatty liver disease. *World Journal of Gastroenterology*, 27(26), 3683- 3696.