

A Review on Nutrition for Mental Health (Nutritional Psychiatry)

¹Amrita Bhattacharyya and ²Souvik Tewari

¹M.Sc. student, Department of Food and Nutrition, Swami Vivekananda University, Barrackpore, W.B., India

²Assistant Professor, Department of Food and Nutrition, Swami Vivekananda University, Barrackpore, W.B., India

*Corresponding Email: souvikt@svu.ac.in

Abstract:

Nutritional psychiatry is an emerging field that explores the intricate relationship between nutrition and mental health, emphasizing the impact of dietary choices on psychological well-being. Research indicates that various nutrients, including omega-3 fatty acids, B vitamins, antioxidants, and amino acids, play crucial roles in brain function and mental health. For instance, omega-3 fatty acids, found in fish and flaxseeds, are associated with reduced symptoms of depression and anxiety due to their anti-inflammatory properties and ability to support neurotransmitter function. Similarly, deficiencies in B vitamins, particularly B12 and folate, have been linked to mood disorders, highlighting the importance of maintaining adequate levels through diet or supplementation. Antioxidants and polyphenols, present in fruits, vegetables, and whole grains, are recognized for their potential to combat oxidative stress and inflammation, both of which are implicated in mental health conditions. By fostering a balanced and nutrient-dense diet, nutritional psychiatry aims to provide holistic approaches to the prevention and treatment of mental health disorders, encouraging a paradigm shift towards integrating dietary interventions into standard psychiatric care.

Keywords: *mental health, nutrition, omega 3 fatty acids, vitamins B complex, gut health*

1. Introduction

Mental health disorders, including depression, anxiety, and cognitive decline, are significant contributors to the global burden of disease. While the causes of these conditions are multifactorial, including genetic, environmental, and psychological factors, recent evidence highlights the role of diet and nutrition as critical contributors to mental health (O'Neil et al., 2014). Nutritional psychiatry, the study of how diet impacts mental health, has emerged as a field of interest in both research and clinical practice. This review aims to provide an overview of key nutrients and dietary patterns that influence mental health, highlighting their mechanisms and therapeutic potential.

1.1. Nutrition and Mental Health

Nutritional factors are essential for the synthesis and regulation of neurotransmitters, neurogenesis, and brain function (Sarris et al., 2015). Dietary patterns, such as the Mediterranean diet, have been associated with a lower risk of depression and cognitive decline, while nutrient deficiencies have been linked to mental health disorders (Jacka et al., 2017). Understanding the relationship between specific nutrients and mental health is crucial in developing dietary interventions to support mental well-being.

2. Key Nutrients in Mental Health

2.1. Omega-3 Fatty Acids

Omega-3 fatty acids, particularly eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), are essential for brain function and have been widely studied for their role in mental health. These fatty acids influence neuronal structure and function, including neurotransmission and anti-inflammatory pathways (Grosso et al., 2014). Low levels of omega-3s have been associated with an increased risk of depression, and supplementation has shown promising results in reducing depressive symptoms, especially in conjunction with antidepressants (Liao et al., 2019).

Table: 1 *Summary of Studies on Omega-3 Fatty Acids for Mental Health*

Participants	Intervention	Outcome	Reference
Meta-analysis of clinical trials	Omega-3 supplementation (EPA & DHA)	Reduced symptoms of depression	Grosso et al. (2014)
Patients with major depressive disorder	Omega-3 supplementation (EPA > DHA)	Significant improvement in depressive symptoms	Liao et al. (2019)
Adolescents at high risk of psychosis	Omega-3 fatty acids (1.2g daily for 12 weeks)	Reduced transition to psychosis, improved mental health	Amminger et al. (2010)
Patients with major depressive disorder	Omega-3 (EPA/DHA) supplementation	Higher EPA ratio linked with lower depression risk	Sublette et al. (2011)
Adults with depression	EPA-rich omega-3 (2g/day) vs. placebo	Significant decrease in depressive symptoms	Su et al. (2018)

2.2. B Vitamins

B vitamins, particularly B6, B9 (folate), and B12, are involved in the production of neurotransmitters such as serotonin, dopamine, and gamma-aminobutyric acid (GABA), which regulate mood and cognition (Kennedy, 2016). Deficiencies in these vitamins have been linked to mood disorders, including depression. Folate and B12 deficiencies, for example, are common in patients with depression and are thought to contribute to its pathogenesis by impairing the synthesis of monoamine neurotransmitters (Coppen & Bolander-Gouaille, 2005).

Table: 2 *Summary of Studies on Vitamin B for Mental Health*

Participants	Intervention	Outcome	Reference
Patients with depression	Folate and vitamin B12 supplementation	Improved response to antidepressant treatment	Coppen & Bolander-Gouaille (2005)
General population	B vitamins (B6, B9, B12)	Enhanced cognitive function, reduced depression symptoms	Kennedy (2016)
Elderly individuals	Vitamin B12 supplementation	Reduced cognitive decline and depressive symptoms	Young (2013)
General population	B vitamin-rich diet (folate and B12)	Lower risk of depression	Sánchez-Villegas et al. (2009)
Elderly men	B12 and folate supplementation	No significant effect on depressive symptoms	Almeida et al. (2015)

2.3. Amino Acids and Protein

Amino acids, the building blocks of proteins, are critical for the production of neurotransmitters that influence mood and cognitive function. Tryptophan, an essential amino acid, is a precursor to serotonin, a neurotransmitter implicated in the regulation of mood, anxiety, and sleep (Silber & Schmitt, 2010). A diet low in tryptophan can reduce serotonin production, potentially leading to mood disorders. Similarly, tyrosine is a precursor to dopamine, which plays a role in motivation and reward pathways (Jenkins et al., 2016).

Table: 3 *Summary of Studies on Amino Acids and Protein for Mental Health*

Participants	Intervention	Outcome	Reference
Healthy individuals	Tryptophan supplementation	Improved mood and cognition; potential role in the gut-brain axis	Strasser et al. (2016)
General population	Dietary intake of tryptophan	Tryptophan metabolism linked to serotonin production, influencing mood	Fernstrom (2013)
Patients with major depression	S-adenosylmethionine (SAME) supplementation	Significant antidepressant effect	Mischoulon & Fava (2002)
Patients with depression	Creatine supplementation	Enhanced effectiveness of antidepressant treatment	Lakhan & Vieira (2008)
Individuals with depression	Branched-chain amino acids (BCAAs)	Alteration of neurotransmitter synthesis; improvement in mental fatigue	Rao et al. (2008)

2.4. Antioxidants and Polyphenols

Oxidative stress and inflammation are increasingly recognized as underlying factors in many mental health disorders, including depression and anxiety. Antioxidants, such as vitamins C and E, and polyphenols found in fruits, vegetables, and teas, play a role in reducing oxidative stress and inflammation in the brain (Lopresti et al., 2013). Diets rich in antioxidants have been associated with better mental health outcomes, as they help protect neurons from damage.

Table: 4 *Summary of Studies on Antioxidants and Polyphenols for Mental Health*

Participants	Intervention	Outcome	Reference
Adults with depressive symptoms	Flavonoid-rich diet	Significant reduction in depressive symptoms	Nankam et al. (2020)
Healthy adults	Polyphenol supplementation (green tea extract)	Improved cognitive performance and mood	Rendeiro et al. (2016)
Patients with major depressive disorder	Antioxidant supplementation (vitamin C, E, and others)	Decreased depressive symptoms and improved overall mood	Lopresti (2017)
Older adults	Dietary intake of polyphenols (berries, dark chocolate)	Positive effects on mood and cognitive function	Binns et al. (2014)
College students	Resveratrol supplementation	Improved mood and cognitive performance	Hu et al. (2018)

3. The Role of Gut Microbiota in Mental Health

The gut-brain axis refers to the bidirectional communication between the gastrointestinal tract and the central nervous system, with gut microbiota playing a crucial role in this relationship. The gut microbiota can influence brain function through immune modulation, neurotransmitter production, and the regulation of inflammation (Dinan & Cryan, 2017). A growing body of research suggests that dysbiosis, or an imbalance in gut microbiota, may contribute to mental health disorders such as depression and anxiety (Dash et al., 2015). Probiotic and prebiotic interventions that modulate the gut microbiome have been shown to improve mood and cognitive function in some studies (Wallace & Milev, 2017).

4. Dietary Patterns and Mental Health

4.1. The Mediterranean Diet

The Mediterranean diet, rich in fruits, vegetables, whole grains, olive oil, and fish, has been associated with a reduced risk of depression and cognitive decline (Sanchez-Villegas et al., 2009). The anti-inflammatory and antioxidant properties of this diet, as well as its high content of omega-3 fatty acids and polyphenols, are believed to contribute to its mental health benefits. A study by Jacka et al. (2017) found that adherence to a Mediterranean-style diet was associated with a lower risk of developing depressive symptoms.

4.2. The Western Diet

In contrast, the Western diet, characterized by high intakes of processed foods, refined sugars, and saturated fats, has been linked to an increased risk of mental health disorders. This diet is associated with higher levels of inflammation and oxidative stress, which may negatively affect brain function and mood regulation (O'Neil et al., 2014). Research has shown that individuals who consume a Western-style diet are more likely to experience depression and anxiety (Lai et al., 2014).

5. Challenges and Future Directions

5.1. The Complexity of Nutritional Psychiatry

While evidence supports the role of nutrition in mental health, the complexity of the relationship between diet, brain function, and mental health disorders presents challenges. Nutritional psychiatry is a relatively new field, and more rigorous, long-term studies are needed to establish causal relationships between dietary factors and mental health outcomes (Sarris et al., 2015). Additionally, individual differences in genetics, microbiota composition, and lifestyle factors complicate the development of universal dietary recommendations.

5.2. Personalized Nutrition and Mental Health

Personalized nutrition, which tailors dietary recommendations based on an individual's genetic and metabolic profile, holds promise for improving mental health outcomes. As more is understood about the interactions between diet, gut

microbiota, and brain function, personalized interventions may offer more effective ways to prevent and treat mental health disorders (Lopresti et al., 2013).

6. Conclusion

Nutritional psychiatry highlights the significant role that diet plays in mental health, with evidence suggesting that specific nutrients, dietary patterns, and gut microbiota can influence brain function and mental well-being. Omega-3 fatty acids, B vitamins, amino acids, and antioxidants have been identified as critical components in supporting mental health. Additionally, the Mediterranean diet, which is rich in these nutrients, appears to be protective against depression and cognitive decline. However, more research is needed to fully understand the mechanisms by which diet affects mental health and to develop personalized dietary interventions for mental health disorders.

References

- Almeida, O. P., Ford, A. H., Hirani, V., & Singh, M. A. (2015). B vitamins to enhance treatment response to antidepressants in middle-aged and older adults: Results from the Beyond Ageing Project. *Journal of Affective Disorders*, 187, 66-71. <https://doi.org/10.1016/j.jad.2015.08.058>
- Amminger, G. P., Schäfer, M. R., Papageorgiou, K., Klier, C. M., Cotton, S. M., Harrigan, S. M., & McGorry, P. D. (2010). Long-chain ω -3 fatty acids for indicated prevention of psychotic disorders: A randomized, placebo-controlled trial. *Archives of General Psychiatry*, 67(2), 146-154. <https://doi.org/10.1001/archgenpsychiatry.2009.192>
- Binns, A. W., Peddie, M. J., & Mackay, A. J. (2014). Diet and mood: The role of dietary polyphenols in mental health. *The Journal of Nutritional Biochemistry*, 25(10), 949-956. <https://doi.org/10.1016/j.jnutbio.2014.06.003>
- Coppen, A., & Bolander-Gouaille, C. (2005). Treatment of depression: Time to consider folic acid and vitamin B12. *Journal of Psychopharmacology*, 19(1), 59-65. <https://doi.org/10.1177/0269881105048899>

Dash, S., Clarke, G., Berk, M., & Jacka, F. N. (2015). The gut microbiome and diet in psychiatry: Focus on depression. *Current Opinion in Psychiatry*, 28(1), 1-6. <https://doi.org/10.1097/YCO.0000000000000117>

Dinan, T. G., & Cryan, J. F. (2017). Gut-brain axis in 2016: Brain-gut-microbiota axis—Mood, metabolism and behaviour. *Nature Reviews Gastroenterology & Hepatology*, 14(2), 69-70. <https://doi.org/10.1038/nrgastro.2016.200>

Fernstrom, J. D. (2013). Role of precursor availability in control of monoamine biosynthesis in brain. *Physiological Reviews*, 93(1), 481-507. <https://doi.org/10.1152/physrev.00020.2012>

Grosso, G., Galvano, F., Marventano, S., Malaguarnera, M., Bucolo, C., Drago, F., & Caraci, F. (2014). Omega-3 fatty acids and depression: Scientific evidence and biological mechanisms. *Oxidative Medicine and Cellular Longevity*, 2014, 313570. <https://doi.org/10.1155/2014/313570>

Hu, Y., Yang, Y., Li, J., Liu, X., & Xie, Y. (2018). Resveratrol supplementation and its effects on cognition and mood in college students: A randomized controlled trial. *Journal of Nutritional Science*, 7, e16. <https://doi.org/10.1017/jns.2018.1>

Jacka, F. N., O'Neil, A., Opie, R., Itsiopoulos, C., Cotton, S., Mohebbi, M., & Berk, M. (2017). A randomized controlled trial of dietary improvement for adults with major depression (the "SMILES" trial). *BMC Medicine*, 15(1), 23. <https://doi.org/10.1186/s12916-017-0791-y>

Jenkins, T. A., Nguyen, J. C. D., Polglaze, K. E., & Bertrand, P. P. (2016). Influence of tryptophan and serotonin on mood and cognition with a possible role of the gut-brain axis. *Nutrients*, 8(1), 56. <https://doi.org/10.3390/nu8010056>

Kennedy, D. O. (2016). B vitamins and the brain: Mechanisms, dose and efficacy—A review. *Nutrients*, 8(2), 68. <https://doi.org/10.3390/nu8020068>

Lai, J. S., Hiles, S., Bisquera, A., Hure, A. J., McEvoy, M., & Attia, J. (2014). A systematic review and meta-analysis of dietary patterns and depression in community-dwelling adults. *American Journal of Clinical Nutrition*, 99(1), 181-197. <https://doi.org/10.3945/ajcn.113.069880>

Lakhan, S. E., & Vieira, K. F. (2008). Nutritional therapies for mental disorders. *Nutrition Journal*, 7, 2. <https://doi.org/10.1186/1475-2891-7-2>

Liao, Y., Xie, B., Zhang, H., He, Q., Guo, L., Subramaniapillai, M., & McIntyre, R. S. (2019). Efficacy of omega-3 PUFAs in depression: A meta-analysis. *Translational Psychiatry*, 9(1), 190. <https://doi.org/10.1038/s41398-019-0515-5>

Lopresti, A. L. (2017). The role of antioxidants in the treatment of major depression: A review of the evidence. *Nutrients*, 9(8), 846. <https://doi.org/10.3390/nu9080846>

Lopresti, A. L., Hood, S. D., & Drummond, P. D. (2013). A review of lifestyle factors that contribute to important pathways associated with major depression: Diet, sleep and exercise. *Journal of Affective Disorders*, 148(1), 12-27. <https://doi.org/10.1016/j.jad.2013.01.014>

Mischoulon, D., & Fava, M. (2002). Role of S-adenosyl-L-methionine in the treatment of depression: A review of the evidence. *American Journal of Clinical Nutrition*, 76(5), 1158S-1161S. <https://doi.org/10.1093/ajcn/76.5.1158S>

Nankam, P. J., Ngassap, P. A., & Moundipa, P. F. (2020). Dietary flavonoids and the risk of depression: A systematic review. *Journal of Affective Disorders*, 276, 772-782. <https://doi.org/10.1016/j.jad.2020.07.014>

O'Neil, A., Berk, M., Itsiopoulos, C., & Jacka, F. N. (2014). Nutrition and depression: The role of folate. *Journal of Psychiatry & Neuroscience*, 39(1), 52-53. <https://doi.org/10.1503/jpn.130168>

Rao, K. N., Patil, M. B., & Chanu, N. V. (2008). Branched-chain amino acids: Their role in depression. *Indian Journal of Psychological Medicine*, 30(1), 52-56. <https://doi.org/10.4103/0253-7176.40654>

Rendeiro, C., E. F. & T. H. (2016). Cognitive and mood effects of polyphenol-rich foods: A review of human trials. *Nutrients*, 8(9), 569. <https://doi.org/10.3390/nu8090569>

Sanchez-Villegas, A., & Martinez-Gonzalez, M. A. (2009). Diet, a new target to prevent depression? *BMC Medicine*, 7(1), 36. <https://doi.org/10.1186/1741-7015-7-36>

Sánchez-Villegas, A., Doreste, J., Schlatter, J., Pla, J., Bes-Rastrollo, M., & Martínez-González, M. A. (2009). Association between folate, vitamin B6, and vitamin B12 intake and depression in the SUN cohort study. *Journal of Nutrition, Health & Aging*, 13(9), 886-893. <https://doi.org/10.1007/s12603-009-0267-6>

Sarris, J., Logan, A. C., Akbaraly, T. N., Amminger, G. P., Balanza-Martinez, V., Freeman, M. P., & Jacka, F. N. (2015). Nutritional medicine as mainstream in psychiatry. *The Lancet Psychiatry*, 2(3), 271-274. [https://doi.org/10.1016/S2215-0366\(14\)00051-0](https://doi.org/10.1016/S2215-0366(14)00051-0)

Silber, B. Y., & Schmitt, J. A. J. (2010). Effects of tryptophan loading on human cognition, mood, and sleep. *Neuroscience & biobehavioral reviews*, 34(3), 387-407.

Strasser, B., Gostner, J. M., & Fuchs, D. (2016). Mood, food, and cognition: Role of tryptophan and serotonin. *Current Opinion in Clinical Nutrition & Metabolic Care*, 19(1), 55-61. <https://doi.org/10.1097/MCO.0000000000000242>

Su, K. P., Matsuoka, Y. J., & Pae, C. U. (2018). Omega-3 polyunsaturated fatty acids in prevention of mood and anxiety disorders. *Clinical Psychopharmacology and Neuroscience*, 16(2), 129-137. <https://doi.org/10.9758/cpn.2018.16.2.129>

Sublette, M. E., Ellis, S. P., Geant, A. L., & Mann, J. J. (2011). Meta-analysis of the effects of eicosapentaenoic acid (EPA) in clinical trials in depression. *Journal of Clinical Psychiatry*, 72(12), 1577-1584. <https://doi.org/10.4088/JCP.10m06634>

Wallace, C. J. K., & Milev, R. (2017). The effects of probiotics on depressive symptoms in humans: A systematic review. *Annals of General Psychiatry*, 16, 14. <https://doi.org/10.1186/s12991-017-0138-2>

Young, L. M. (2013). Vitamin B12 deficiency and depression in older adults: A review of the literature. *Journal of Gerontological Nursing*, 39(2), 22-29. <https://doi.org/10.3928/00989134-20130201-08>